



Zinc Fact Sheet: Zinc Fertilizer in the Environment

June 2015

Executive Summary

- Zinc is an essential plant nutrient, but if added to soil in excessive amounts may cause concern from an environmental viewpoint.
- Loadings of zinc to soil from fertilizer use are small in comparison to situations where animal manures or sewage biosolids are applied to agricultural land.
- Most concern regarding applications of zinc to soil is therefore related to disposal/re-use of animal/human wastes, or where zinc is added to soil from industrial/consumer sources, e.g. runoff from galvanized structures.
- To our knowledge, there are no instances reported where zinc added to soil via fertilizers has caused environmental harm.
- A simple mass balance model was developed to examine potential scenarios for how zinc application in fertilizers causes accumulation of zinc in soil, and how this relates to critical toxicity concentrations for zinc in soil.
- The model was parameterized for a typical agricultural soil scenario and assumed farmers keep adding fertilizer Zn even though available Zn in soil increases to sufficiency levels (an unlikely scenario). The time for Zn to accumulate in soil to concentrations approaching toxicity thresholds was more than a millennium, and even then, removal in produce and through leaching could keep soil concentrations below critical toxicity thresholds ad infinitum.
- Data from a long-term field trial in Virginia on a neutral soil, where high rates of Zn fertilizer were applied to corn over a 20-year period (rates equivalent to more than 500 years of normal rates of Zn application), indicated no adverse effects on crop growth.
- The environmental risk of Zn in fertilizers is therefore extremely low.

Background

Zinc (Zn) is an essential element for both crops and humans, and many soils around the world are potentially Zn deficient (Alloway 2008)(Figure 1).

Zinc is one of the key factors affecting human health in developing countries and UNICEF has a program of Zn supplementation of child diets to overcome these deficiencies either directly by dietary intervention (adding Zn compound powders directly to foods) or through biofortification (enhancing Zn concentrations in food crops by breeding and/or fertilization) (Brown et al. 2009; Hotz 2009).

Deficiency of Zn in soil can therefore adversely affect not only crop production through reduced yields, but also affect the health of humans eating food produced from those soils.

Hence many farmers now require (and request) Zn-enriched fertilizers for use in crop production.




Figure 1. Map of zinc-deficient agricultural soils throughout the world. (Alloway, 2008)

Additions of Zinc to Soils in Fertilizers

Rates of Zn addition to soil through fertilization vary from <1 kg Zn/ha up to 25 kg Zn/ha depending on the crop type and soil and environmental conditions (Alloway 2008). In very deficient soils these rates may need to be re-applied for several years until the available Zn status in soil has increased, and then rates of Zn application can be reduced, or the frequency of Zn application reduced.

A typical Zn fertilizer application to soil would be an addition of ~2.5 kg Zn/ha. For comparison, the addition of 5 tonnes of animal manure or sewage biosolids to soil could easily add 1.5–10.0 kg Zn/ha or even more (Benke et al. 2008).

What is the Fate of Fertilizer Zinc Added to Soil?

Fertilizer Zn added to soil does not diffuse far from the point of application (Hettiarachchi et al. 2008), as most soils bind Zn very strongly and this reduces movement of fertilizer Zn (Figure 2) from the granule (or fluid injection point). Binding of Zn in soil is predominantly governed by soil pH (Sauve et al. 2000; Tiller and Hodgson 1962), with alkaline soils binding Zn more strongly (hence alkaline soils are more prone to be Zn deficient and require Zn fertilization).




Figure 2. Map of Zn concentrations in soil around a Zn-enriched fertilizer granule 5 weeks after application to soil. Reprinted by Permission, ASA, CSSA, SSSA.

Zinc that diffuses into the soil can either be taken up by crop plants (and the Zn removed in produce) or may be lost from soil by leaching - leaching losses are usually greater in sandy, low pH soils which generally are not Zn deficient and therefore are not fertilized with Zn. Losses of Zn through runoff will generally be small as Zn binds so strongly to soil particles (Brümmer et al. 1983). Any Zn added in excess of crop removals and leaching will accumulate over time

in soil building up the total Zn concentrations and the available Zn concentrations. Once available Zn is built up to a specific level (dependent on soil and crop type and climatic conditions), further addition of Zn will not be required for several years (Brennan 2001) as the soil becomes “non-responsive” to Zn fertilizer.

Zinc Toxicity in Soil

If Zn accumulates to high enough concentrations in soil, it may cause toxicity to plants or to soil microorganisms (Chaney 1993). This phenomenon was first observed in soils contaminated with runoff from galvanized structures (Millikan 1947) and is most commonly observed only in contaminated urban/industrial soils or where wastes have been applied to land, e.g. biosolids (Chaudri et al. 1993).

Much research has been conducted to set “safe” levels for Zn in soil, and naturally these are not single values across all soils but vary, due to variable ability of soil to bind Zn, depending on soil pH and other soil properties (e.g. cation exchange capacity (CEC)) (Smolders et al. 2004; Smolders et al. 2009; Warne et al. 2008).

Could Fertilizer Zinc Cause Toxicity in Soils?

It is highly unlikely that fertilizer zinc could cause toxicity in soils for several reasons:

1. Amounts of Zn added to soil in fertilizers are very small;
2. Zinc is removed in agricultural produce and as Zn concentrations in soil increase, so too does the Zn concentrations in produce removed from the land (which is good for human and animal nutrition); and
3. Fertilizer Zn has residual value; i.e. once Zn levels in soil have been increased through repeated application of Zn fertilizers, there is no need for farmers to continually add more Zn to soils and applications are often reduced in frequency or rate to match crop removals.

A simple model of fertilizer Zn behavior in soil was developed to model inputs in fertilizer, crop offtake and losses through leaching. This model also calculates the maximum concentration in soil likely to lead to toxicity to compare to soil concentrations resultant from fertilizer addition. Note the model does not include the likely reduction in Zn inputs once Zn concentrations in soils have increased to sufficiency levels (point 3 above). Normally farmers would not apply Zn if the extractable Zn concentrations in soil reach sufficiency levels.



Naturally, accumulation of Zn depends on fertilizer management regime, crop type, soil type, etc. Results for a typical corn cropping system in mid-western USA are presented in Figure 3.

It is evident from Figure 3 that, even assuming farmers do not reduce the rate or frequency of Zn applications as Zn concentrations and availability in soil increase over time (which they normally would), it takes many centuries for Zn concentrations in soil to increase significantly, and soil Zn concentrations do not exceed toxicity thresholds at “equilibrium” under this scenario. Other scenarios can be assessed using the simple Excel model.

The modelling results are confirmed by results from a long-term field trial where up to 900 kg Zn/ha was cumulatively added to a neutral (pH 7.0) silty clay soil over a period of 19 years (Payne et al. 1988). Yields of corn and silage were unaffected even though soil Zn concentrations reached >250 mg/kg. Movement of Zn to depth was minimal also. Given that normal rates of Zn application are ~1 kg/ha/yr, this trial confirms that in the long term (500-1000 years) there is a low risk of fertiliser Zn posing environmental problems.




Figure 3. Predicted increase in Zn concentrations in soil in relation to the concentration expected to cause toxicity. Assumptions for each yearly cycle are: addition of fertilizer Zn of 2 kg/ha, atmospheric deposition of 1 g Zn/ha, corn yield of 10 t/ha, and deep drainage of 300 mm; and for the soil properties: soil pH 6.0, CEC 15 cmolc/kg (sandy loam soil), and initial total soil Zn concentration of 40 mg/kg. Partitioning of Zn between soil and soil solution according to the relationship determined by (Degryse et al. 2009)⁹, toxicity thresholds for plants determined according to soil pH and CEC¹⁸ and toxicity thresholds for nitrifying bacteria determined according to soil pH.⁴

References

1. **Alloway, BJ.** *Zinc in Soils and Crop Nutrition*. International Zinc Association and International Fertilizer Association, Brussels, Belgium. 2008.
2. **Benke MB, Indraratne SR, Hao KY, Chang C, Goh TB.** "Trace element changes in soil after long-term cattle manure applications." *Journal of Environmental Quality*. 37:798-807. 2008.
3. **Brennan R.** "Residual value of zinc fertiliser for production of wheat." *Australian Journal of Experimental Agriculture*. 41:541-547. 2001.
4. **Broos K, Warne MSJ, Heemsbergen DA, Stevens DP, Barnes MB, Correll RL, McLaughlin MJ.** Soil factors controlling the toxicity of Cu and Zn to microbial processes in Australian soils. *Environmental Toxicology and Chemistry* 26:583-590. 2007.
5. **Brown KH, Engle-Stone R, Krebs NF, Peerson JM.** "Dietary intervention strategies to enhance zinc nutrition: Promotion and support of breast feeding for infants and young children." *Food and Nutrition Bulletin*. 30:S144-S171. 2009.
6. **Brümmer G, Tiller KT, Herms U, Clayton P.** "Absorption-desorption and/or precipitation-dissolution processes of zinc in soils." *Geoderma*. 31:337-354. 1983.
7. **Chaney RL.** "Zinc phytotoxicity." *Zinc in Soils and Plants*. Ed. AD Robson. Kluwer Academic Publishers, Dordrecht. 1993.
8. **Chaudri AM, McGrath SP, Giller KE, Rietz E, Sauerbeck DR.** "Enumeration of indigenous rhizobium leguminosarum biovar trifolii in soils previously treated with metal-contaminated sewage sludge." *Soil Biology and Biochemistry*. 25:301-309. 1993.
9. **Degryse F, Smolders E, Parker DR.** Partitioning of trace metals in soils: Concepts, methodologies, predictions and applications - a review. *Europe J Soil Sci* 60:590-612. 2009.
10. **Hettiarachchi GM, McLaughlin MJ, Schekel KG, Chittleborough DJ, Newville M, Sutton S, Lombi E.** "Evidence for different reaction pathways for liquid- and granular-micronutrients in a calcareous soil." *Soil Science Society of America Journal*. 72:98-110. 2008.
11. **Hotz C.** "The potential to improve zinc status through biofortification of staple food crops with zinc." *Food and Nutrition Bulletin*. 30:S172-S178. 2009.
12. **Millikan CR.** "Zinc toxicity in flax grown in a wire netting 'bird cage'." *Journal of the Australian Institute of Agricultural Science*. 13:64-67. 1947.
13. **Payne GG, Martens DC, Winarko C, Perera NF.** "Form and availability of copper and zinc following long-term copper sulfate and zinc sulfate applications." *Journal of Environmental Quality*. 17:707-711. 1988.
14. **Sauve S, Hendershot W, Allen HE.** "Solid-solution partitioning of metals in contaminated soils: Dependence on pH, total metal burden, and organic matter." *Environmental Science and Technology*. 34:1125-1131. 2000.
15. **Smolders E, Buekers J, Oliver I, McLaughlin MJ.** "Soil properties affecting toxicity of zinc to soil microbial properties in laboratory-spiked and field-contaminated soils." *Environmental Toxicology and Chemistry*. 23:2633-2640. 2004.
16. **Smolders E, Oorts K, Van Sprang P, Schoeters I, Janssen CJ, McGrath SP, McLaughlin MJ.** "Toxicity of trace metals in soil as affected by soil type and aging after contamination: Using calibrated bioavailability models to set ecological soil standards." *Environmental Toxicology and Chemistry*. 28:1633-1642. 2009.
17. **Tiller KG, Hodgson JF.** "The specific sorption of cobalt and zinc by layer silicates." *Clay Mineralogy*. 9:393-403. 1962.
18. **Warne MSJ, Heemsbergen D, McLaughlin M, Bell M, Broos K, Whatmuff M, Barry G, Nash D, Pritchard D, Penney N.** "Models for the field-based toxicity of copper and zinc salts to wheat in 11 Australian soils and comparison to laboratory-based models." *Environmental Pollution*. 156:707-714 2008.

Authored By:

Prof. Mike McLaughlin
Dr. Fien Degryse
Fertilizer Technology Research Centre
Adelaide University
School of Agriculture Food and Wine



The Zinc Nutrient Initiative was launched by The International Zinc Association (IZA) in response to the critical issue of zinc deficiency in soils, crops and humans. To learn more, visit: www.zinc.org/crops.

